

22. Orthogonal Polynomials

URS W. HOCHSTRASSER¹

Contents

	Page
Mathematical Properties	773
22.1. Definition of Orthogonal Polynomials	773
22.2. Orthogonality Relations	774
22.3. Explicit Expressions	775
22.4. Special Values	777
22.5. Interrelations	777
22.6. Differential Equations	781
22.7. Recurrence Relations	782
22.8. Differential Relations	783
22.9. Generating Functions	783
22.10. Integral Representations	784
22.11. Rodrigues' Formula	785
22.12. Sum Formulas	785
22.13. Integrals Involving Orthogonal Polynomials	785
22.14. Inequalities	786
22.15. Limit Relations	787
22.16. Zeros	787
22.17. Orthogonal Polynomials of a Discrete Variable	788
Numerical Methods	788
22.18. Use and Extension of the Tables	788
22.19. Least Square Approximations	790
22.20. Economization of Series	791
References	792
Table 22.1. Coefficients for the Jacobi Polynomials $P_n^{(\alpha, \beta)}(x)$	793
$n=0(1)6$	
Table 22.2. Coefficients for the Ultraspherical Polynomials $C_n^{(\alpha)}(x)$ and for x^n in Terms of $C_m^{(\alpha)}(x)$	794
$n=0(1)6$	
Table 22.3. Coefficients for the Chebyshev Polynomials $T_n(x)$ and for x^n in Terms of $T_m(x)$	795
$n=0(1)12$	
Table 22.4. Values of the Chebyshev Polynomials $T_n(x)$	795
$n=0(1)12, x=.2(.2)1, 10D$	
Table 22.5. Coefficients for the Chebyshev Polynomials $U_n(x)$ and for x^n in Terms of $U_m(x)$	796
$n=0(1)12$	
Table 22.6. Values of the Chebyshev Polynomials $U_n(x)$	796
$n=0(1)12, x=.2(.2)1, 10D$	

¹ Guest Worker, National Bureau of Standards, from The American University. (Presently, Atomic Energy Commission, Switzerland.)

	Page
Table 22.7. Coefficients for the Chebyshev Polynomials $C_n(x)$ and for x^n in Terms of $C_m(x)$ $n=0(1)12$	797
Table 22.8. Coefficients for the Chebyshev Polynomials $S_n(x)$ and for x^n in Terms of $S_m(x)$ $n=0(1)12$	797
Table 22.9. Coefficients for the Legendre Polynomials $P_n(x)$ and for x^n in Terms of $P_m(x)$ $n=0(1)12$	798
Table 22.10. Coefficients for the Laguerre Polynomials $L_n(x)$ and for x^n in Terms of $L_m(x)$ $n=0(1)12$	799
Table 22.11. Values of the Laguerre Polynomials $L_n(x)$ $n=0(1)12, x=.5, 1, 3, 5, 10, \text{ Exact or } 10D$	800
Table 22.12. Coefficients for the Hermite Polynomials $H_n(x)$ and for x^n in Terms of $H_m(x)$ $n=0(1)12$	801
Table 22.13. Values of the Hermite Polynomials $H_n(x)$ $n=0(1)12, x=.5, 1, 3, 5, 10, \text{ Exact or } 11S$	802

22. Orthogonal Polynomials

Mathematical Properties

22.1. Definition of Orthogonal Polynomials

A system of polynomials $f_n(x)$, degree $[f_n(x)] = n$, is called orthogonal on the interval $a \leq x \leq b$, with respect to the weight function $w(x)$, if

22.1.1

$$\int_a^b w(x) f_n(x) f_m(x) dx = 0 \quad (n \neq m; n, m = 0, 1, 2, \dots)$$

The weight function $w(x)[w(x) \geq 0]$ determines the system $f_n(x)$ up to a constant factor in each polynomial. The specification of these factors is referred to as standardization. For suitably standardized orthogonal polynomials we set

22.1.2

$$\int_a^b w(x) f_n^2(x) dx = h_n, \quad f_n(x) = k_n x^n + k'_n x^{n-1} + \dots \quad (n = 0, 1, 2, \dots)$$

These polynomials satisfy a number of relationships of the same general form. The most important ones are:

Differential Equation

$$22.1.3 \quad g_2(x) f_n'' + g_1(x) f_n' + a_n f_n = 0$$

where $g_2(x)$, $g_1(x)$ are independent of n and a_n a constant depending only on n .

Recurrence Relation

$$22.1.4 \quad f_{n+1} = (a_n + x b_n) f_n - c_n f_{n-1}$$

where

22.1.5

$$b_n = \frac{k_{n+1}}{k_n}, \quad a_n = b_n \left(\frac{k'_{n+1}}{k_{n+1}} - \frac{k'_n}{k_n} \right), \quad c_n = \frac{k_{n+1} k_{n-1} h_n}{k_n^2 h_{n-1}}$$

Rodrigues' Formula

$$22.1.6 \quad f_n = \frac{1}{e_n w(x)} \frac{d^n}{dx^n} \{ w(x) [g(x)]^n \}$$

where $g(x)$ is a polynomial in x independent of n . The system $\left\{ \frac{df_n}{dx} \right\}$ consists again of orthogonal polynomials.




FIGURE 22.1. *Jacobi Polynomials* $P_n^{(\alpha, \beta)}(x)$, $\alpha = 1.5$, $\beta = .5$, $n = 1(1)5$.

22.2. Orthogonality Relations

	$f_n(x)$	Name of Polynomial	a	b	$w(x)$	Standardization	h_n	Remarks
22.2.1	$P_n^{(\alpha, \beta)}(x)$	Jacobi	-1	1	$(1-x)^\alpha(1+x)^\beta$	$P_n^{(\alpha, \beta)}(1) = \binom{n+\alpha}{n}$	$\frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{n!\Gamma(n+\alpha+\beta+1)}$	$\alpha > -1, \beta > -1$
22.2.2	$G_n(p, q, x)$	Jacobi	0	1	$(1-x)^{p-\frac{1}{2}}x^{q-\frac{1}{2}}$	$k_n = 1$	$\frac{n!\Gamma(n+q)\Gamma(n+p)\Gamma(n+p-q+1)}{(2n+p)\Gamma^2(2n+p)}$	$p-q > -1, q > 0$
22.2.3	$C_n^{(\alpha)}(x)$	Ultraspherical (Gegenbauer)	-1	1	$(1-x^2)^{\alpha-\frac{1}{2}}$	$C_n^{(\alpha)}(1) = \binom{n+2\alpha-1}{n}$ $(\alpha \neq 0)$	$\frac{\pi^{21-2\alpha}\Gamma(n+2\alpha)}{n!(n+\alpha)[\Gamma(\alpha)]^2} \quad \alpha \neq 0$	$\alpha > -\frac{1}{2}$
						$C_n^{(0)}(1) = \frac{2}{n},$ $C_0^{(0)}(1) = 1$	$\frac{2\pi}{n^2} \quad \alpha = 0$	
22.2.4	$T_n(x)$	Chebyshev of the first kind	-1	1	$(1-x^2)^{-\frac{1}{2}}$	$T_n(1) = 1$	$\begin{cases} \frac{\pi}{2} & n \neq 0 \\ \pi & n = 0 \end{cases}$	
22.2.5	$U_n(x)$	Chebyshev of the second kind	-1	1	$(1-x^2)^{\frac{1}{2}}$	$U_n(1) = n+1$	$\frac{\pi}{2}$	∞
22.2.6	$C_n(x)$	Chebyshev of the first kind	-2	2	$\left(1-\frac{x^2}{4}\right)^{-\frac{1}{2}}$	$C_n(2) = 2$	$\begin{cases} 4\pi & n \neq 0 \\ 8\pi & n = 0 \end{cases}$	
22.2.7	$S_n(x)$	Chebyshev of the second kind	-2	2	$\left(1-\frac{x^2}{4}\right)^{\frac{1}{2}}$	$S_n(2) = n+1$	π	
22.2.8	$T_n^*(x)$	Shifted Chebyshev of the first kind	0	1	$(x-x^2)^{-\frac{1}{2}}$	$T_n^*(1) = 1$	$\begin{cases} \frac{\pi}{2} & n \neq 0 \\ \pi & n = 0 \end{cases}$	
22.2.9	$U_n^*(x)$	Shifted Chebyshev of the second kind	0	1	$(x-x^2)^{\frac{1}{2}}$	$U_n^*(1) = n+1$	$\frac{\pi}{8} \quad *$	
22.2.10	$P_n(x)$	Legendre (Spherical)	-1	1	1	$P_n(1) = 1$	$\frac{2}{2n+1}$	
22.2.11	$P_n^*(x)$	Shifted Legendre	0	1	1		$\frac{1}{2n+1}$	

*See page 11.

22.2. Orthogonality Relations—Continued

	22.2.12	$L_n^{(\alpha)}(x)$	Generalized Laguerre	0	∞	$e^{-x}x^\alpha$	$k_n = \frac{(-1)^n}{n!}$	$\frac{\Gamma(\alpha+n+1)}{n!}$	$\alpha > -1$
	22.2.13	$L_n(x)$	Laguerre	0	∞	e^{-x}	$k_n = \frac{(-1)^n}{n!}$	1	
*	22.2.14	$H_n(x)$	Hermite	$-\infty$	∞	e^{-x^2}	$e_n = (-1)^n$	$\sqrt{\pi}2^n n!$	
*	22.2.15	$He_n(x)$	Hermite	$-\infty$	∞	$e^{-\frac{x^2}{2}}$	$e_n = (-1)^n$	$\sqrt{2\pi}n!$	

* See page II.

22.3. Explicit Expressions

$$f_n(x) = d_n \sum_{m=0}^N c_m g_m(x)$$

	$f_n(x)$	N	d_n	c_m	$g_m(x)$	k_n	Remarks
22.3.1	$P_n^{(\alpha, \beta)}(x)$	n	$\frac{1}{2^n}$	$\binom{n+\alpha}{m} \binom{n+\beta}{n-m}$	$(x-1)^{n-m}(x+1)^m$	$\frac{1}{2^n} \binom{2n+\alpha+\beta}{n}$	$\alpha > -1, \beta > -1$
22.3.2	$P_n^{(\alpha, \beta)}(x)$	n	$\frac{\Gamma(\alpha+n+1)}{n! \Gamma(\alpha+\beta+n+1)}$	$\binom{n}{m} \frac{\Gamma(\alpha+\beta+n+m+1)}{2^m \Gamma(\alpha+m+1)}$	$(x-1)^m$	$\frac{1}{2^n} \binom{2n+\alpha+\beta}{n}$	$\alpha > -1, \beta > -1$
22.3.3	$G_n(p, q, x)$	n	$\frac{\Gamma(q+n)}{\Gamma(p+2n)}$	$(-1)^m \binom{n}{m} \frac{\Gamma(p+2n-m)}{\Gamma(q+n-m)}$	x^{n-m}	1	$p-q > -1, q > 0$
22.3.4	$C_n^{(\alpha)}(x)$	$\left[\frac{n}{2}\right]$	$\frac{1}{\Gamma(\alpha)}$	$(-1)^m \frac{\Gamma(\alpha+n-m)}{m!(n-2m)!}$	$(2x)^{n-2m}$	$\frac{2^n}{n!} \frac{\Gamma(\alpha+n)}{\Gamma(\alpha)}$	$\alpha > -\frac{1}{2}, \alpha \neq 0$
22.3.5	$C_n^{(0)}(x)$	$\left[\frac{n}{2}\right]$	1	$(-1)^m \frac{(n-m-1)!}{m!(n-2m)!}$	$(2x)^{n-2m}$	$\frac{2^n}{n} \quad n \neq 0$	$n \neq 0, C_0^{(0)}(1) = 1$
22.3.6	$T_n(x)$	$\left[\frac{n}{2}\right]$	$\frac{n}{2}$	$(-1)^m \frac{(n-m-1)!}{m!(n-2m)!}$	$(2x)^{n-2m}$	2^{n-1}	
22.3.7	$U_n(x)$	$\left[\frac{n}{2}\right]$	1	$(-1)^m \frac{(n-m)!}{m!(n-2m)!}$	$(2x)^{n-2m}$	2^n	
22.3.8	$P_n(x)$	$\left[\frac{n}{2}\right]$	$\frac{1}{2^n}$	$(-1)^m \binom{n}{m} \binom{2n-2m}{n}$	x^{n-2m}	$\frac{(2n)!}{2^n (n!)^2}$	
22.3.9	$L_n^{(\alpha)}(x)$	n	1	$(-1)^m \binom{n+\alpha}{n-m} \frac{1}{m!}$	x^m	$\frac{(-1)^n}{n!}$	$\alpha > -1$
22.3.10	$H_n(x)$	$\left[\frac{n}{2}\right]$	$n!$	$(-1)^m \frac{1}{m!(n-2m)!}$	$(2x)^{n-2m}$	2^n	see 22.11
22.3.11	$He_n(x)$	$\left[\frac{n}{2}\right]$	$n!$	$(-1)^m \frac{1}{m! 2^m (n-2m)!}$	x^{n-2m}	1	




FIGURE 22.2. *Jacobi Polynomials* $P_n^{(\alpha, \beta)}(x)$,
 $\alpha=1(.2)2$, $\beta=-.5$, $n=5$.




FIGURE 22.3. *Jacobi Polynomials* $P_n^{(\alpha, \beta)}(x)$,
 $\alpha=1.5$, $\beta=-.8(.2)0$, $n=5$.

Explicit Expressions Involving Trigonometric Functions

$$f_n(\cos \theta) = \sum_{m=0}^n a_m \cos(n-2m)\theta$$

	$f_n(\cos \theta)$	a_m	Remarks
22.3.12	$C_n^{(\alpha)}(\cos \theta)$	$\frac{\Gamma(\alpha+m)\Gamma(\alpha+n-m)}{m!(n-m)![\Gamma(\alpha)]^2}$	$\alpha \neq 0$
22.3.13	$P_n(\cos \theta)$	$\frac{1}{4^n} \binom{2m}{m} \binom{2n-2m}{n-m}$	

$$22.3.14 \quad C_n^{(0)}(\cos \theta) = \frac{2}{n} \cos n\theta$$

$$22.3.15 \quad T_n(\cos \theta) = \cos n\theta$$

$$22.3.16 \quad U_n(\cos \theta) = \frac{\sin(n+1)\theta}{\sin \theta}$$





FIGURE 22.4. *Gegenbauer (Ultraspherical) Polynomials* $C_n^{(\alpha)}(x)$, $\alpha=.5$, $n=2(1)5$.

22.4. Special Values

	$f_n(x)$	$f_n(-x)$	$f_n(1)$	$f_n(0)$	$f_0(x)$	$f_1(x)$
22.4.1	$P_n^{(\alpha, \beta)}(x)$	$(-1)^n P_n^{(\beta, \alpha)}(x)$	$\binom{n+\alpha}{n}^*$		1	$\frac{1}{2}[\alpha - \beta + (\alpha + \beta + 2)x]$
22.4.2	$C_n^{(\alpha)}(x)$ $\alpha \neq 0$	$(-1)^n C_n^{(\alpha)}(x)$	$\binom{n+2\alpha-1}{n}$	$\begin{cases} 0, n=2m+1 \\ (-1)^{n/2} \frac{\Gamma(\alpha+n/2)}{\Gamma(\alpha)(n/2)!}, n=2m \end{cases}$	1	$2\alpha x$
22.4.3	$C_n^{(0)}(x)$	$(-1)^n C_n^{(0)}(x)$	$\frac{2}{n}, n \neq 0$	$\begin{cases} \frac{(-1)^m}{m}, n=2m \neq 0 \\ 0, n=2m+1 \end{cases}$	1	$2x$
22.4.4	$T_n(x)$	$(-1)^n T_n(x)$	1	$\begin{cases} (-1)^m, n=2m \\ 0, n=2m+1 \end{cases}$	1	x
22.4.5	$U_n(x)$	$(-1)^n U_n(x)$	$n+1$	$\begin{cases} (-1)^m, n=2m \\ 0, n=2m+1 \end{cases}$	1	$2x$
22.4.6	$P_n(x)$	$(-1)^n P_n(x)$	1	$\begin{cases} \frac{(-1)^m}{4^m} \binom{2m}{m}, n=2m^* \\ 0, n=2m+1 \end{cases}$	1	x
22.4.7	$L_n^{(\alpha)}(x)$			$\binom{n+\alpha}{n}$	1	$-x + \alpha + 1$
22.4.8	$H_n(x)$	$(-1)^n H_n(x)$		$\begin{cases} (-1)^m \frac{(2m)!}{m!}, n=2m \\ 0, n=2m+1 \end{cases}$	1	$2x$


 FIGURE 22.5. Gegenbauer (Ultraspherical) Polynomials $C_n^{(\alpha)}(x)$, $\alpha = .2(.2)1$, $n = 5$.

22.5. Interrelations

Interrelations Between Orthogonal Polynomials of the Same Family

Jacobi Polynomials

22.5.1

$$P_n^{(\alpha, \beta)}(x) = \frac{\Gamma(2n + \alpha + \beta + 1)}{n! \Gamma(n + \alpha + \beta + 1)} G_n\left(\alpha + \beta + 1, \beta + 1, \frac{x+1}{2}\right)$$

22.5.2

$$G_n(p, q, x) = \frac{n! \Gamma(n+p)}{\Gamma(2n+p)} P_n^{(p-q, q-1)}(2x-1)$$

(see [22.21]).

22.5.3

$$F_n(p, q, x) = (-1)^n n! \frac{\Gamma(q)}{\Gamma(q+n)} P_n^{(p-q, q-1)}(2x-1)$$

(see [22.13]).

Ultraspherical Polynomials

$$22.5.4 \quad C_n^{(0)}(x) = \lim_{\alpha \rightarrow 0} \frac{1}{\alpha} C_n^{(\alpha)}(x)$$

Chebyshev Polynomials

$$22.5.5 \quad T_n(x) = \frac{1}{2} C_n(2x) = T_n^*\left(\frac{1+x}{2}\right)$$

$$22.5.6 \quad T_n(x) = U_n(x) - x U_{n-1}(x)$$

$$22.5.7 \quad T_n(x) = xU_{n-1}(x) - U_{n-2}(x)$$

$$22.5.8 \quad T_n(x) = \frac{1}{2} [U_n(x) - U_{n-2}(x)]$$

$$22.5.9 \quad U_n(x) = S_n(2x) = U_n^* \left(\frac{1+x}{2} \right)$$

$$22.5.10 \quad U_{n-1}(x) = \frac{1}{1-x^2} [xT_n(x) - T_{n+1}(x)]$$

$$22.5.11 \quad C_n(x) = 2T_n \left(\frac{x}{2} \right) = 2T_n^* \left(\frac{x+2}{4} \right)$$

$$22.5.12 \quad C_n(x) = S_n(x) - S_{n-2}(x)$$

$$22.5.13 \quad S_n(x) = U_n \left(\frac{x}{2} \right) = U_n^* \left(\frac{x+2}{4} \right)$$

$$22.5.14 \quad T_n^*(x) = T_n(2x-1) = \frac{1}{2} C_n(4x-2)$$

(see [22.22]).

$$22.5.15 \quad U_n^*(x) = S_n(4x-2) = U_n(2x-1)$$

(see [22.22]).

Generalized Laguerre Polynomials

$$22.5.16 \quad L_n^{(0)}(x) = L_n(x)$$

$$22.5.17 \quad L_n^{(m)}(x) = (-1)^m \frac{d^m}{dx^m} [L_{n+m}(x)]$$

Hermite Polynomials

$$22.5.18 \quad He_n(x) = 2^{-n/2} H_n \left(\frac{x}{\sqrt{2}} \right)$$

(see [22.20]).

$$22.5.19 \quad H_n(x) = 2^{n/2} He_n(x\sqrt{2})$$

(see [22.13], [22.20]).

Interrelations Between Orthogonal Polynomials of Different Families

Jacobi Polynomials

22.5.20

$$P_n^{(\alpha-\frac{1}{2}, \alpha-\frac{1}{2})}(x) = \frac{\Gamma(2\alpha)\Gamma(\alpha+n+\frac{1}{2})}{\Gamma(2\alpha+n)\Gamma(\alpha+\frac{1}{2})} C_n^{(\alpha)}(x)$$

22.5.21

$$P_n^{(\alpha, \frac{1}{2})}(x) = \frac{(\frac{1}{2})_{n+1}}{\sqrt{\frac{x+1}{2}} (\alpha+\frac{1}{2})_{n+1}} C_{2n+1}^{(\alpha+\frac{1}{2})} \left(\sqrt{\frac{x+1}{2}} \right)$$

$$22.5.22 \quad P_n^{(\alpha, -\frac{1}{2})}(x) = \frac{(\frac{1}{2})_n}{(\alpha+\frac{1}{2})_n} C_{2n}^{(\alpha+\frac{1}{2})} \left(\sqrt{\frac{x+1}{2}} \right)$$

$$22.5.23 \quad P_n^{(-\frac{1}{2}, -\frac{1}{2})}(x) = \frac{1}{4^n} \binom{2n}{n} T_n(x)$$

$$22.5.24 \quad P_n^{(0,0)}(x) = P_n(x)$$

Ultraspherical Polynomials

22.5.25

$$C_{2n}^{(\alpha)}(x) = \frac{\Gamma(\alpha+n)n!2^{2n}}{\Gamma(\alpha)(2n)!} P_n^{(\alpha-\frac{1}{2}, -\frac{1}{2})}(2x^2-1) \quad (\alpha \neq 0)$$

22.5.26

$$C_{2n+1}^{(\alpha)}(x) = \frac{\Gamma(\alpha+n+1)n!2^{2n+1}}{\Gamma(\alpha)(2n+1)!} x P_n^{(\alpha-\frac{1}{2}, \frac{1}{2})}(2x^2-1) \quad (\alpha \neq 0)$$

22.5.27

$$C_n^{(\alpha)}(x) = \frac{\Gamma(\alpha+\frac{1}{2})\Gamma(2\alpha+n)}{\Gamma(2\alpha)\Gamma(\alpha+n+\frac{1}{2})} P_n^{(\alpha-\frac{1}{2}, \alpha-\frac{1}{2})}(x) \quad (\alpha \neq 0)$$

22.5.28

$$C_n^{(0)}(x) = \frac{2}{n} T_n(x) = 2 \frac{(n-1)!}{\Gamma(n+\frac{1}{2})} \sqrt{\pi} P_n^{(-\frac{1}{2}, -\frac{1}{2})}(x) \quad *$$

Chebyshev Polynomials

$$22.5.29 \quad T_{2n+1}(x) = \frac{n! \sqrt{\pi}}{\Gamma(n+\frac{1}{2})} x P_n^{(-\frac{1}{2}, \frac{1}{2})}(2x^2-1)$$

$$22.5.30 \quad U_{2n}(x) = \frac{n! \sqrt{\pi}}{\Gamma(n+\frac{1}{2})} P_n^{(\frac{1}{2}, -\frac{1}{2})}(2x^2-1)$$

$$22.5.31 \quad T_n(x) = \frac{n! \sqrt{\pi}}{\Gamma(n+\frac{1}{2})} P_n^{(-\frac{1}{2}, -\frac{1}{2})}(x)$$

$$22.5.32 \quad U_n(x) = \frac{(n+1)! \sqrt{\pi}}{2\Gamma(n+\frac{3}{2})} P_n^{(\frac{1}{2}, \frac{1}{2})}(x)$$




FIGURE 22.6. Chebyshev Polynomials $T_n(x)$, $n=1(1)5$.

*See page II.




FIGURE 22.7. Chebyshev Polynomials $U_n(x)$, $n=1(1)5$.

$$22.5.33 \quad T_n(x) = \frac{n}{2} C_n^{(0)}(x)$$

$$22.5.34 \quad U_n(x) = C_n^{(1)}(x)$$

Legendre Polynomials

$$22.5.35 \quad P_n(x) = P_n^{(0,0)}(x)$$

$$22.5.36 \quad P_n(x) = C_n^{(1/2)}(x)$$

$$22.5.37$$

$$\frac{d^m}{dx^m} [P_n(x)] = 1 \cdot 3 \cdots (2m-1) C_{n-m}^{(m+1/2)}(x) \quad (m \leq n)$$

Generalized Laguerre Polynomials

$$22.5.38 \quad L_n^{(-1/2)}(x) = \frac{(-1)^n}{n! 2^{2n}} H_{2n}(\sqrt{x})$$

$$22.5.39 \quad L_n^{(1/2)}(x) = \frac{(-1)^n}{n! 2^{2n+1} \sqrt{x}} H_{2n+1}(\sqrt{x})$$

Hermite Polynomials

$$22.5.40 \quad H_{2m}(x) = (-1)^m 2^{2m} m! L_m^{(-1/2)}(x^2)$$

$$22.5.41 \quad H_{2m+1}(x) = (-1)^m 2^{2m+1} m! x L_m^{(1/2)}(x^2)$$

Orthogonal Polynomials as Hypergeometric Functions (see chapter 15)

$$f_n(x) = {}_dF(a, b; c; g(x))$$

For each of the listed polynomials there are numerous other representations in terms of hypergeometric functions.

	$f_n(x)$	d	a	b	c	$g(x)$
22.5.42	$P_n^{(\alpha, \beta)}(x)$	$\binom{n+\alpha}{n}$	$-n$	$n+\alpha+\beta+1$	$\alpha+1$	$\frac{1-x}{2}$
22.5.43	$P_n^{(\alpha, \beta)}(x)$	$\binom{2n+\alpha+\beta}{n} \left(\frac{x-1}{2}\right)^n$	$-n$	$-n-\alpha$	$-2n-\alpha-\beta$	$\frac{2}{1-x}$
22.5.44	$P_n^{(\alpha, \beta)}(x)$	$\binom{n+\alpha}{n} \left(\frac{1+x}{2}\right)^n$	$-n$	$-n-\beta$	$\alpha+1$	$\frac{x-1}{x+1}$
22.5.45	$P_n^{(\alpha, \beta)}(x)$	$\binom{n+\beta}{n} \left(\frac{x-1}{2}\right)^n$	$-n$	$-n-\alpha$	$\beta+1$	$\frac{x+1}{x-1}$
22.5.46	$C_n^{(\alpha)}(x)$	$\frac{\Gamma(n+2\alpha)}{n! \Gamma(2\alpha)}$	$-n$	$n+2\alpha$	$\alpha+\frac{1}{2}$	$\frac{1-x}{2}$
22.5.47	$T_n(x)$	1	$-n$	n	$\frac{1}{2}$	$\frac{1-x}{2}$
22.5.48	$U_n(x)$	$n+1$	$-n$	$n+2$	$\frac{3}{2}$	$\frac{1-x}{2}$
22.5.49	$P_n(x)$	1	$-n$	$n+1$	1	$\frac{1-x}{2}$
22.5.50	$P_n(x)$	$\binom{2n}{n} \left(\frac{x-1}{2}\right)^n$	$-n$	$-n$	$-2n$	$\frac{2}{1-x}$
22.5.51	$P_n(x)$	$\binom{2n}{n} \left(\frac{x}{2}\right)^n$	$-\frac{n}{2}$	$\frac{1-n}{2}$	$\frac{1}{2}-n$	$\frac{1}{x^2}$
22.5.52	$P_{2n}(x)$	$(-1)^n \frac{(2n)!}{2^{2n} (n!)^2}$	$-n$	$n+\frac{1}{2}$	$\frac{1}{2}$	x^2
22.5.53	$P_{2n+1}(x)$	$(-1)^n \frac{(2n+1)!}{2^{2n} (n!)^2} x$	$-n$	$n+\frac{3}{2}$	$\frac{3}{2}$	x^2

*See page II.

Orthogonal Polynomials as Confluent Hypergeometric Functions (see chapter 13)

$$22.5.54 \quad L_n^{(\alpha)}(x) = \binom{n+\alpha}{n} M(-n, \alpha+1, x)$$

Orthogonal Polynomials as Parabolic Cylinder Functions (see chapter 19)

$$22.5.55 \quad H_n(x) = 2^n U\left(\frac{1}{2} - \frac{1}{2}n, \frac{3}{2}, x^2\right)$$

$$22.5.56 \quad H_{2m}(x) = (-1)^m \frac{(2m)!}{m!} M\left(-m, \frac{1}{2}, x^2\right)$$

22.5.57

$$\bullet \quad H_{2m+1}(x) = (-1)^m \frac{(2m+1)!}{m!} 2x M\left(-m, \frac{3}{2}, x^2\right)$$




FIGURE 22.8. Legendre Polynomials $P_n(x)$, $n=2(1)5$.

22.5.58

$$H_n(x) = 2^{n/2} e^{x^2/2} D_n(\sqrt{2}x) = 2^{n/2} e^{x^2/2} U\left(-n - \frac{1}{2}, \sqrt{2}x\right)$$

$$22.5.59 \quad He_n(x) = e^{x^2/4} D_n(x) = e^{x^2/4} U\left(-n - \frac{1}{2}, x\right)$$

Orthogonal Polynomials as Legendre Functions (see chapter 8)

22.5.60

$$C_n^{(\alpha)}(x) =$$

$$\frac{\Gamma(\alpha + \frac{1}{2}) \Gamma(2\alpha + n)}{n! \Gamma(2\alpha)} \left[\frac{1}{4}(x^2 - 1)\right]^{\frac{1}{2} - \alpha} P_{n+\alpha-\frac{1}{2}}^{(\frac{1}{2}-\alpha)}(x) \quad (\alpha \neq 0)$$




FIGURE 22.9. Laguerre Polynomials $L_n(x)$, $n=2(1)5$.




FIGURE 22.10. Hermite Polynomials $\frac{H_n(x)}{n!}$, $n=2(1)5$.

22.6. Differential Equations

$$g_2(x)y'' + g_1(x)y' + g_0(x)y = 0$$

	y	$g_2(x)$	$g_1(x)$	$g_0(x)$
22.6.1	$P_n^{(\alpha, \beta)}(x)$	$1-x^2$	$\beta - \alpha - (\alpha + \beta + 2)x$	$n(n + \alpha + \beta + 1)$
22.6.2	$(1-x)^\alpha(1+x)^\beta P_n^{(\alpha, \beta)}(x)$	$1-x^2$	$\alpha - \beta + (\alpha + \beta - 2)x$	$(n+1)(n + \alpha + \beta)$
22.6.3	$(1-x)^{\frac{\alpha+1}{2}}(1+x)^{\frac{\beta+1}{2}} P_n^{(\alpha, \beta)}(x)$	1	0	$\frac{1}{4} \frac{1-\alpha^2}{(1-x)^2} + \frac{1}{4} \frac{1-\beta^2}{(1+x)^2}$ $+ \frac{2n(n + \alpha + \beta + 1) + (\alpha+1)(\beta+1)}{2(1-x^2)}$
22.6.4	$\left(\sin \frac{x}{2}\right)^{\alpha+\frac{1}{2}} \left(\cos \frac{x}{2}\right)^{\beta+\frac{1}{2}} P_n^{(\alpha, \beta)}(\cos x)$	1	0	$\frac{1-4\alpha^2}{16 \sin^2 \frac{x}{2}} + \frac{1-4\beta^2}{16 \cos^2 \frac{x}{2}}$ $+ \left(n + \frac{\alpha + \beta + 1}{2}\right)^2$
22.6.5	$C_n^{(\alpha)}(x)$	$1-x^2$	$-(2\alpha+1)x$	$n(n+2\alpha)$
22.6.6	$(1-x^2)^{\alpha-\frac{1}{2}} C_n^{(\alpha)}(x)$	$1-x^2$	$(2\alpha-3)x$	$(n+1)(n+2\alpha-1)$
22.6.7	$(1-x^2)^{\frac{\alpha}{2}-\frac{1}{4}} C_n^{(\alpha)}(x)$	1	0	$\frac{(n+\alpha)^2}{1-x^2} + \frac{2+4\alpha-4\alpha^2+x^2}{4(1-x^2)^2}$
22.6.8	$(\sin x)^\alpha C_n^{(\alpha)}(\cos x)$	1	0	$(n+\alpha)^2 + \frac{\alpha(1-\alpha)}{\sin^2 x}$
22.6.9	$T_n(x)$	$1-x^2$	$-x$	n^2
22.6.10	$T_n(\cos x)$	1	0	n^2
22.6.11	$\frac{1}{\sqrt{1-x^2}} T_n(x); U_{n-1}(x)$	*	$1-x^2$	$-3x$
22.6.12	$U_n(x)$	$1-x^2$	$-3x$	$n(n+2)$
22.6.13	$P_n(x)$	$1-x^2$	$-2x$	$n(n+1)$
22.6.14	$\sqrt{1-x^2} P_n(x)$	1	0	$\frac{n(n+1)}{1-x^2} + \frac{1}{(1-x^2)^2}$
22.6.15	$L_n^{(\alpha)}(x)$	x	$\alpha+1-x$	n
22.6.16	$e^{-x} x^{\alpha/2} L_n^{(\alpha)}(x)$	*	x	$2+1$
22.6.17	$e^{-x/2} x^{(\alpha+1)/2} L_n^{(\alpha)}(x)$	1	0	$n + \frac{\alpha}{2} + 1 - \frac{\alpha^2}{4x}$
22.6.18	$e^{-x^2/2} x^{\alpha+\frac{1}{2}} L_n^{(\alpha)}(x^2)$	1	0	$\frac{2n+\alpha+1}{2x} + \frac{1-\alpha^2}{4x^2} - \frac{1}{4}$
22.6.19	$H_n(x)$	1	$-2x$	$4n+2\alpha+2-x^2 + \frac{1-4\alpha^2}{4x^2}$
22.6.20	$e^{-\frac{x^2}{2}} H_n(x)$	1	0	$2n+1-x^2$
22.6.21	$He_n(x)$	1	$-x$	n

*See page II.

22.7. Recurrence Relations

Recurrence Relations With Respect to the Degree n

$$a_{1n}f_{n+1}(x) = (a_{2n} + a_{3n}x)f_n(x) - a_{4n}f_{n-1}(x)$$

	f_n	a_{1n}	a_{2n}	a_{3n}	a_{4n}
22.7.1	$P_n^{(\alpha, \beta)}(x)$	$2(n+1)(n+\alpha+\beta+1)$ $(2n+\alpha+\beta)$	$(2n+\alpha+\beta+1)(\alpha^2-\beta^2)$	$(2n+\alpha+\beta)_2$	$2(n+\alpha)(n+\beta)$ $(2n+\alpha+\beta+2)$
22.7.2	$G_n(p, q, x)$	$(2n+p-2)_2(2n+p-1)$	$-[2n(n+p)+q(p-1)]$ $(2n+p-2)_2$	$(2n+p-2)_4$ $(2n+p-1)$	$n(n+q-1)(n+p-1)$ $(n+p-q)(2n+p+1)$
22.7.3	$C_n^{(\alpha)}(x)$	$n+1$	0	$2(n+\alpha)$	$n+2\alpha-1$
22.7.4	$T_n(x)$	1	0	2	1
22.7.5	$U_n(x)$	1	0	2	1
22.7.6	$S_n(x)$	1	0	1	1
22.7.7	$C_n(x)$	1	0	1	1
22.7.8	$T_n^*(x)$	1	-2	4	1
22.7.9	$U_n^*(x)$	1	-2	4	1
22.7.10	$P_n(x)$	$n+1$	0	$2n+1$	n
22.7.11	$P_n^*(x)$	$n+1$	$-2n-1$	$4n+2$	n
22.7.12	$L_n^{(\alpha)}(x)$	$n+1$	$2n+\alpha+1$	-1	$n+\alpha$
22.7.13	$H_n(x)$	1	0	2	$2n$
22.7.14	$He_n(x)$	1	0	1	n

Miscellaneous Recurrence Relations

Jacobi Polynomials

22.7.15

$$\left(n + \frac{\alpha}{2} + \frac{\beta}{2} + 1\right)(1-x)P_n^{(\alpha+1, \beta)}(x) \\ = (n+\alpha+1)P_n^{(\alpha, \beta)}(x) - (n+1)P_{n+1}^{(\alpha, \beta)}(x)$$

22.7.16

$$\left(n + \frac{\alpha}{2} + \frac{\beta}{2} + 1\right)(1+x)P_n^{(\alpha, \beta+1)}(x) \\ = (n+\beta+1)P_n^{(\alpha, \beta)}(x) + (n+1)P_{n+1}^{(\alpha, \beta)}(x)$$

22.7.17

$$(1-x)P_n^{(\alpha+1, \beta)}(x) + (1+x)P_n^{(\alpha, \beta+1)}(x) = 2P_n^{(\alpha, \beta)}(x)$$

22.7.18

$$(2n+\alpha+\beta)P_n^{(\alpha-1, \beta)}(x) = (n+\alpha+\beta)P_n^{(\alpha, \beta)}(x) \\ - (n+\beta)P_{n-1}^{(\alpha, \beta)}(x)$$

22.7.19

$$(2n+\alpha+\beta)P_n^{(\alpha, \beta-1)}(x) = (n+\alpha+\beta)P_n^{(\alpha, \beta)}(x) \\ + (n+\alpha)P_{n-1}^{(\alpha, \beta)}(x)$$

$$22.7.20 \quad P_n^{(\alpha, \beta-1)}(x) - P_n^{(\alpha-1, \beta)}(x) = P_{n-1}^{(\alpha, \beta)}(x)$$

Ultraspherical Polynomials

22.7.21

$$2\alpha(1-x^2)C_{n-1}^{(\alpha+1)}(x) = (2\alpha+n-1)C_{n-1}^{(\alpha)}(x) - nx C_n^{(\alpha)}(x)$$

22.7.22

$$= (n+2\alpha)x C_n^{(\alpha)}(x) \\ - (n+1)C_{n+1}^{(\alpha)}(x)$$

$$22.7.23 \quad (n+\alpha)C_{n+1}^{(\alpha-1)}(x) = (\alpha-1)[C_{n+1}^{(\alpha)}(x) - C_{n-1}^{(\alpha)}(x)]$$

Chebyshev Polynomials

22.7.24

$$2T_m(x)T_n(x) = T_{n+m}(x) + T_{n-m}(x) \quad (n \geq m) \quad *$$

22.7.25

$$2(x^2-1)U_{m-1}(x)U_{n-1}(x) = T_{n+m}(x) - T_{n-m}(x) \quad (n \geq m)$$

22.7.26

$$2T_m(x)U_{n-1}(x) = U_{n+m-1}(x) + U_{n-m-1}(x) \quad (n > m)$$

22.7.27

$$2T_n(x)U_{m-1}(x) = U_{n+m-1}(x) - U_{n-m-1}(x) \quad (n > m)$$

$$22.7.28 \quad 2T_n(x)U_{n-1}(x) = U_{2n-1}(x)$$

Generalized Laguerre Polynomials

22.7.29

$$L_n^{(\alpha+1)}(x) = \frac{1}{x} [(x-n)L_n^{(\alpha)}(x) + (\alpha+n)L_{n-1}^{(\alpha)}(x)]$$

22.7.30

$$L_n^{(\alpha-1)}(x) = L_n^{(\alpha)}(x) - L_{n-1}^{(\alpha)}(x)$$

22.7.31

$$L_n^{(\alpha+1)}(x) = \frac{1}{x} [(n+\alpha+1)L_n^{(\alpha)}(x) - (n+1)L_{n+1}^{(\alpha)}(x)]$$

22.7.32

$$L_n^{(\alpha-1)}(x) = \frac{1}{n+\alpha} [(n+1)L_{n+1}^{(\alpha)}(x) - (n+1-x)L_n^{(\alpha)}(x)]$$

22.8. Differential Relations

$$g_1(x) \frac{d}{dx} f_n(x) = g_1(x) f_n(x) + g_0(x) f_{n-1}(x)$$

	f_n	g_1	g_1	g_0
22.8.1	$P_n^{(\alpha, \beta)}(x)$	$(2n+\alpha+\beta)(1-x^2)$	$n[\alpha-\beta-(2n+\alpha+\beta)x]$	$2(n+\alpha)(n+\beta)$
22.8.2	$C_n^{(\alpha)}(x)$	$1-x^2$	$-nx$	$n+2\alpha-1$
22.8.3	$T_n(x)$	$1-x^2$	$-nx$	n
22.8.4	$U_n(x)$	$1-x^2$	$-nx$	$n+1$
22.8.5	$P_n(x)$	$1-x^2$	$-nx$	n
22.8.6	$L_n^{(\alpha)}(x)$	x	n	$-(n+\alpha)$
22.8.7	$H_n(x)$	1	0	$2n$
22.8.8	$He_n(x)$	1	0	n

22.9. Generating Functions

$$g(x, z) = \sum_{n=0}^{\infty} a_n f_n(x) z^n$$

$$R = \sqrt{1-2xz+z^2}$$

	$f_n(x)$	a_n	$g(x, z)$	Remarks
22.9.1	$P_n^{(\alpha, \beta)}(x)$	$2^{-\alpha-\beta}$	$R^{-1}(1-z+R)^{-\alpha}(1+z+R)^{-\beta}$	$ z < 1$
22.9.2	$C_n^{(\alpha)}(x)$	$\frac{2^{1-\alpha} \Gamma(\alpha + \frac{1}{2} + n) \Gamma(2\alpha)}{\Gamma(\alpha + \frac{1}{2}) \Gamma(2\alpha + n)}$	$R^{-1}(1-xz+R)^{1-\alpha}$	$ z < 1, \alpha \neq 0$
22.9.3	$C_n^{(\alpha)}(x)$	1	$R^{-2\alpha}$	$ z < 1, \alpha \neq 0$
22.9.4	$C_n^{(0)}(x)$	1	$-\ln R^2$	$ z < 1$
22.9.5	$C_n^{(\alpha)}(x)$	$\frac{\Gamma(2\alpha)}{\Gamma(\alpha + \frac{1}{2}) \Gamma(2\alpha + n)}$	$e^{i \cos \theta} \left(\frac{z}{2} \sin \theta \right)^{\frac{1}{2}-\alpha} J_{\alpha-\frac{1}{2}}(z \sin \theta)$	$x = \cos \theta$
22.9.6	$T_n(x)$	2	$\left(\frac{1-x^2}{R^2} + 1 \right)$	$-1 < x < 1, z < 1$
22.9.7	$T_n(x)$	$\frac{\sqrt{2}}{4^n} \binom{2n}{n}$	$R^{-1}(1-xz+R)^{1/2}$	$-1 < x < 1, z < 1$
22.9.8	$T_n(x)$	$\frac{1}{n}$	$1 - \frac{1}{2} \ln R^2$	$a_0 = 1, -1 < x < 1, z < 1$
22.9.9	$T_n(x)$	1	$\frac{1-xz}{R^2}$	$-1 < x < 1, z < 1$
22.9.10	$U_n(x)$	1	R^{-2}	$-1 < x < 1, z < 1$
22.9.11	$U_n(x)$	$\frac{\sqrt{2}}{4^{n+1}} \binom{2n+2}{n+1}$	$\frac{1}{R} (1-xz+R)^{-1/2}$	$-1 < x < 1, z < 1$

*See page II.

22.9. Generating Functions—Continued

$$g(x, z) = \sum_{n=0}^{\infty} a_n f_n(x) z^n \quad R = \sqrt{1-2xz+z^2}$$

	$f_n(x)$	a_n	$g(x, z)$	Remarks
22.9.12	$P_n(x)$	1	R^{-1}	$-1 < x < 1$ $ z < 1$
22.9.13	$P_n(x)$	$\frac{1}{n!}$	$e^{xz \cos \theta} J_0(z \sin \theta)$	$x = \cos \theta$
22.9.14	$S_n(x)$	1	$(1-xz+z^2)^{-1}$	$-2 < x < 2$ $ z < 1$
22.9.15	$L_n^{(\alpha)}(x)$	1	$(1-z)^{-\alpha-1} \exp\left(\frac{xz}{z-1}\right)$	$ z < 1$
22.9.16	$L_n^{(\alpha)}(x)$	$\frac{1}{\Gamma(n+\alpha+1)}$	$(xz)^{-\frac{1}{2}\alpha} e^{xz} J_n[2(xz)^{1/2}]$	
22.9.17	$H_n(x)$	$\frac{1}{n!}$	e^{2xz-x^2}	
22.9.18	$H_{2n}(x)$	$\frac{(-1)^n}{(2n)!}$	$e^{xz} \cos(2x\sqrt{z})$ *	
22.9.19	$H_{2n+1}(x)$	$\frac{(-1)^n}{(2n+1)!}$	$z^{-1/2} e^{xz} \sin(2x\sqrt{z})$ *	

22.10. Integral Representations

Contour Integral Representations

$f_n(x) = \frac{g_0(x)}{2\pi i} \int_C [g_1(z, x)]^n g_2(z, x) dz$ where C is a closed contour taken around $z=a$ in the positive sense

	$f_n(x)$	$g_0(x)$	$g_1(z, x)$	$g_2(z, x)$	a	Remarks
22.10.1	$P_n^{(\alpha, \beta)}(x)$	$\frac{1}{(1-x)^\alpha(1+x)^\beta}$	$\frac{z^2-1}{2(z-x)}$	$\frac{(1-z)^\alpha(1+z)^\beta}{z-x}$	x	± 1 outside C
22.10.2	$C_n^{(\alpha)}(x)$	1	$1/z$	$(1-2xz+z^2)^{-\alpha} z^{-1}$	0	Both zeros of $1-2xz+z^2$ outside C , $\alpha > 0$
22.10.3	$T_n(x)$	$1/2$	$1/z$	$\frac{1-z^2}{z(1-2xz+z^2)}$	0	Both zeros of $1-2xz+z^2$ outside C
22.10.4	$U_n(x)$	1	$1/z$	$\frac{1}{z(1-2xz+z^2)}$	0	Both zeros of $1-2xz+z^2$ outside C
22.10.5	$P_n(x)$	1	$1/z$	$\frac{1}{z} (1-2xz+z^2)^{-1/2}$	0	Both zeros of $1-2xz+z^2$ outside C
22.10.6	$P_n(x)$	$\frac{1}{2^n}$	$\frac{z^2-1}{z-x}$	$\frac{1}{z-x}$	x	
22.10.7	$L_n^{(\alpha)}(x)$	$e^{2x} x^{-\alpha}$	$\frac{z}{z-x}$	$\frac{z^\alpha}{z-x} e^{-z}$	x	Zero outside C
22.10.8	$L_n^{(\alpha)}(x)$	1	$1 + \frac{x}{z}$	$e^{-z} \left(1 + \frac{z}{x}\right)^n 1/z$	0	$z = -x$ outside C
22.10.9	$H_n(x)$	$n!$	$1/z$	$\frac{e^{2xz-x^2}}{z}$	0	

Miscellaneous Integral Representations

$$22.10.10 \quad C_n^{(\alpha)}(x) = \frac{2^{(1-2\alpha)} \Gamma(n+2\alpha)}{n! [\Gamma(\alpha)]^2} \int_0^\pi [x + \sqrt{x^2-1} \cos \phi]^n (\sin \phi)^{2\alpha-1} d\phi \quad (\alpha > 0)$$

$$22.10.11 \quad C_n^{(\alpha)}(\cos \theta) = \frac{2^{1-\alpha} \Gamma(n+2\alpha)}{n! [\Gamma(\alpha)]^2} (\sin \theta)^{1-2\alpha} \int_0^\theta \frac{\cos(n+\alpha)\phi}{(\cos \phi - \cos \theta)^{1-\alpha}} d\phi \quad (\alpha > 0)$$

*See page II.

$$22.10.12 \quad P_n(\cos \theta) = \frac{1}{\pi} \int_0^\pi (\cos \theta + i \sin \theta \cos \phi)^n d\phi$$

$$22.10.13 \quad P_n(\cos \theta) = \frac{\sqrt{2}}{\pi} \int_0^\pi \frac{\sin(n + \frac{1}{2})\phi d\phi}{(\cos \theta - \cos \phi)^{\frac{1}{2}}}$$

$$22.10.14 \quad L_n^{(\alpha)}(x) = \frac{e^x x^{-\frac{\alpha}{2}}}{n!} \int_0^\infty e^{-t} t^{n+\frac{\alpha}{2}} J_\alpha(2\sqrt{tx}) dt$$

22.10.15

$$H_n(x) = e^{x^2} \frac{2^{n+1}}{\sqrt{\pi}} \int_0^\infty e^{-t^2} t^n \cos\left(2xt - \frac{n}{2}\pi\right) dt$$

22.11. Rodrigues' Formula

$$f_n(x) = \frac{1}{a_n \rho(x)} \frac{d^n}{dx^n} \{ \rho(x) (g(x))^n \}$$

The polynomials given in the following table are the only orthogonal polynomials which satisfy this formula.

	$f_n(x)$	a_n	$\rho(x)$	$g(x)$
22.11.1	$P_n^{(\alpha, \beta)}(x)$	$(-1)^n 2^n n!$	$(1-x)^\alpha (1+x)^\beta$	$1-x^2$
22.11.2	$C_n^{(\alpha)}(x)$	$(-1)^n 2^n n! \frac{\Gamma(2\alpha)\Gamma(\alpha+n+\frac{1}{2})}{\Gamma(\alpha+\frac{1}{2})\Gamma(n+2\alpha)}$	$(1-x^2)^{\alpha-\frac{1}{2}}$	$1-x^2$
22.11.3	$T_n(x)$	$(-1)^n 2^n \frac{\Gamma(n+\frac{1}{2})}{\sqrt{\pi}}$	$(1-x^2)^{-\frac{1}{2}}$	$1-x^2$
22.11.4	$U_n(x)$	$(-1)^n 2^{n+1} \frac{\Gamma(n+\frac{1}{2})}{(n+1)\sqrt{\pi}}$	$(1-x^2)^{\frac{1}{2}}$	$1-x^2$
22.11.5	$P_n(x)$	$(-1)^n 2^n n!$	1	$1-x^2$
22.11.6	$L_n^{(\alpha)}(x)$	$n!$	$e^{-x} x^\alpha$	x
22.11.7	$H_n(x)$	$(-1)^n$	e^{-x^2}	1
22.11.8	$He_n(x)$	$(-1)^n$	$e^{-x^2/2}$	1

22.12. Sum Formulas Christoffel-Darboux Formula

22.12.1

$$\sum_{m=0}^n \frac{1}{h_m} f_m(x) f_m(y) = \frac{k_n}{k_{n+1} h_n} \frac{f_{n+1}(x) f_n(y) - f_n(x) f_{n+1}(y)}{x-y}$$

Miscellaneous Sum Formulas (Only a Limited Selection Is Given Here.)

$$22.12.2 \quad \sum_{m=0}^n T_{2m}(x) = \frac{1}{2} [1 + U_{2n}(x)]$$

$$22.12.3 \quad \sum_{m=0}^{n-1} T_{2m+1}(x) = \frac{1}{2} U_{2n-1}(x)$$

$$22.12.4 \quad \sum_{m=0}^n U_{2m}(x) = \frac{1 - T_{2n+2}(x)}{2(1-x^2)}$$

$$22.12.5 \quad \sum_{m=0}^{n-1} U_{2m+1}(x) = \frac{x - T_{2n+1}(x)}{2(1-x^2)}$$

$$22.12.6 \quad \sum_{m=0}^n L_m^{(\alpha)}(x) L_m^{(\beta)}(y) = L_n^{(\alpha+\beta+1)}(x+y)$$

$$22.12.7 \quad \sum_{m=0}^n \binom{n+\alpha}{m} \mu^{n-m} (1-\mu)^m L_{n-m}^{(\alpha)}(x) = L_n^{(\alpha)}(\mu x)$$

22.12.8

$$H_n(x+y) = \frac{1}{2^{n/2}} \sum_{k=0}^n \binom{n}{k} H_k(\sqrt{2}x) H_{n-k}(\sqrt{2}y)$$

22.13. Integrals Involving Orthogonal Polynomials

22.13.1

$$2n \int_0^x (1-y)^\alpha (1+y)^\beta P_n^{(\alpha, \beta)}(y) dy = P_{n-1}^{(\alpha+1, \beta+1)}(0) - (1-x)^{\alpha+1} (1+x)^{\beta+1} P_{n-1}^{(\alpha+1, \beta+1)}(x)$$

22.13.2

$$\frac{n(2\alpha+n)}{2\alpha} \int_0^x (1-y^2)^{\alpha-\frac{1}{2}} C_n^{(\alpha)}(y) dy = C_{n-1}^{(\alpha+1)}(0) - (1-x^2)^{\alpha+\frac{1}{2}} C_{n-1}^{(\alpha+1)}(x)$$

$$22.13.3 \quad \int_{-1}^1 \frac{T_n(y) dy}{(y-x)\sqrt{1-y^2}} = \pi U_{n-1}(x)$$

$$22.13.4 \quad \int_{-1}^1 \frac{\sqrt{1-y^2} U_{n-1}(y) dy}{(y-x)} = -\pi T_n(x) \quad *$$

$$22.13.5 \quad \int_{-1}^1 (1-x)^{-1/2} P_n(x) dx = \frac{2^{3/2}}{2n+1} \quad *$$

$$22.13.6 \quad \int_0^\pi P_{2n}(\cos \theta) d\theta = \frac{\pi}{16^n} \binom{2n}{n}^2$$

$$22.13.7 \quad \int_0^\pi P_{2n+1}(\cos \theta) \cos \theta d\theta = \frac{\pi}{4^{2n+1}} \binom{2n}{n} \binom{2n+2}{n+1}$$

*See page 11.

22.13.8

$$\int_0^1 x^\lambda P_{2n}(x) dx = \frac{(-1)^n \Gamma\left(n - \frac{\lambda}{2}\right) \Gamma\left(\frac{1}{2} + \frac{\lambda}{2}\right)}{2 \Gamma\left(-\frac{\lambda}{2}\right) \Gamma\left(n + \frac{3}{2} + \frac{\lambda}{2}\right)} \quad (\lambda > -1)$$

22.13.9

$$\int_0^1 x^\lambda P_{2n+1}(x) dx = \frac{(-1)^n \Gamma\left(n + \frac{1}{2} - \frac{\lambda}{2}\right) \Gamma\left(1 + \frac{\lambda}{2}\right)}{2 \Gamma\left(n + 2 + \frac{\lambda}{2}\right) \Gamma\left(\frac{1}{2} - \frac{\lambda}{2}\right)} \quad (\lambda > -2)$$

22.13.10

$$\int_{-1}^z \frac{P_n(t) dt}{\sqrt{x-t}} = \frac{1}{(n+\frac{1}{2})\sqrt{1+x}} [T_n(x) + T_{n+1}(x)]$$

22.13.11

$$\int_z^1 \frac{P_n(t) dt}{\sqrt{t-x}} = \frac{1}{(n+\frac{1}{2})\sqrt{1-x}} [T_n(x) - T_{n+1}(x)]$$

$$22.13.12 \quad \int_z^\infty e^{-t} L_n^{(\alpha)}(t) dt = e^{-z} [L_n^{(\alpha)}(x) - L_{n-1}^{(\alpha)}(x)]$$

22.13.13

$$\Gamma(\alpha + \beta + n + 1) \int_0^z (x-t)^{\beta-1} t^\alpha L_n^{(\alpha)}(t) dt = \Gamma(\alpha + n + 1) \Gamma(\beta) x^{\alpha+\beta} L_n^{(\alpha+\beta)}(x) \quad (\Re \alpha > -1, \Re \beta > 0)$$

22.13.14

$$\int_0^z L_m(t) L_n(x-t) dt = \int_0^z L_{m+n}(t) dt = L_{m+n}(x) - L_{m+n+1}(x)$$

$$22.13.15 \quad \int_0^z e^{-t^2} H_n(t) dt = H_{n-1}(0) - e^{-z^2} H_{n-1}(x)$$

$$22.13.16 \quad \int_0^z H_n(t) dt = \frac{1}{2(n+1)} [H_{n+1}(x) - H_{n+1}(0)]$$

$$22.13.17 \quad \int_{-\infty}^\infty e^{-t^2} H_{2m}(tx) dt = \sqrt{\pi} \frac{(2m)!}{m!} (x^2 - 1)^m$$

22.13.18

$$\int_{-\infty}^\infty e^{-t^2} t H_{2m+1}(tx) dt = \sqrt{\pi} \frac{(2m+1)!}{m!} x (x^2 - 1)^m$$

$$22.13.19 \quad \int_{-\infty}^\infty e^{-t^2} t^n H_n(xt) dt = \sqrt{\pi} n! P_n(x)$$

22.13.20

$$\int_0^\infty e^{-t^2} [H_n(t)]^2 \cos(xt) dt = \sqrt{\pi} 2^{n-1} n! e^{-\frac{x^2}{2}} L_n\left(\frac{x^2}{2}\right)$$

22.14. Inequalities

22.14.1

$$|P_n^{(\alpha, \beta)}(x)| \leq \begin{cases} \binom{n+q}{n} \approx n^q, & \text{if } q = \max(\alpha, \beta) \geq -1/2 \\ & (\alpha > -1, \beta > -1) \\ |P_n^{(\alpha, \beta)}(x')| \approx \sqrt{\frac{1}{n}}, & \text{if } q < -\frac{1}{2} \end{cases}$$

$$x' = \text{maximum point nearest to } \frac{\beta - \alpha}{\alpha + \beta + 1}$$

22.14.2

$$|C_n^{(\alpha)}(x)| \leq \begin{cases} \binom{n+2\alpha-1}{n} & (\alpha > 0) \\ |C_n^{(\alpha)}(x')| & \left(-\frac{1}{2} < \alpha < 0\right) \end{cases}$$

$$x' = 0 \text{ if } n = 2m; x' = \text{maximum point nearest zero} \text{ if } n = 2m + 1$$

22.14.3

$$|C_n^{(\alpha)}(\cos \theta)| < 2^{1-\alpha} \frac{n^{\alpha-1}}{(\sin \theta)^\alpha \Gamma(\alpha)} \quad (0 < \alpha < 1, 0 < \theta < \pi)$$

$$22.14.4 \quad |T_n(x)| \leq 1 \quad (-1 \leq x \leq 1)$$

$$22.14.5 \quad \left| \frac{dT_n(x)}{dx} \right| \leq n^2 \quad (-1 \leq x \leq 1)$$

$$22.14.6 \quad |U_n(x)| \leq n+1 \quad (-1 \leq x \leq 1)$$

$$22.14.7 \quad |P_n(x)| \leq 1 \quad (-1 \leq x \leq 1)$$

$$22.14.8 \quad \left| \frac{dP_n(x)}{dx} \right| \leq \frac{1}{2} n(n+1) \quad (-1 \leq x \leq 1)$$

$$22.14.9 \quad |P_n(x)| \leq \sqrt{\frac{2}{\pi n}} \frac{1}{\sqrt{1-x^2}} \quad (-1 < x \leq 1)^*$$

22.14.10

$$P_n^2(x) - P_{n-1}(x)P_{n+1}(x) < \frac{2n+1}{3n(n+1)} \quad (-1 \leq x \leq 1)$$

22.14.11

$$P_n^2(x) - P_{n-1}(x)P_{n+1}(x) \geq \frac{1 - P_n^2(x)}{(2n-1)(n+1)} \quad (-1 \leq x \leq 1)$$

$$22.14.12 \quad |L_n(x)| \leq e^{x/2} \quad (x \geq 0)$$

$$22.14.13 \quad |L_n^{(\alpha)}(x)| \leq \frac{\Gamma(\alpha + n + 1)}{n! \Gamma(\alpha + 1)} e^{x/2} \quad (\alpha \geq 0, x \geq 0)$$

22.14.14

$$|L_n^{(\alpha)}(x)| \leq \left[2 - \frac{\Gamma(\alpha + n + 1)}{n! \Gamma(\alpha + 1)} \right] e^{x/2} \quad (-1 < \alpha < 0, x \geq 0)$$

*See page 11.

$$22.14.15 \quad |H_{2m}(x)| \leq e^{x^2/2} 2^{2m} m! \left[2 - \frac{1}{2^{2m}} \binom{2m}{m} \right]$$

$$22.14.16 \quad |H_{2m+1}(x)| \leq x e^{x^2/2} \frac{(2m+2)!}{(m+1)!} \quad (x \geq 0)$$

$$22.14.17 \quad |H_n(x)| < e^{x^2/2} k 2^{n/2} \sqrt{n!} \quad k \approx 1.086435$$

22.15. Limit Relations

22.15.1

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n^\alpha} P_n^{(\alpha, \beta)} \left(\cos \frac{x}{n} \right) \right]$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n^\alpha} P_n^{(\alpha, \beta)} \left(1 - \frac{x^2}{2n^2} \right) = \left(\frac{2}{x} \right)^\alpha J_\alpha(x)$$

$$22.15.2 \quad \lim_{n \rightarrow \infty} \left[\frac{1}{n^\alpha} L_n^{(\alpha)} \left(\frac{x}{n} \right) \right] = x^{-\alpha/2} J_\alpha(2\sqrt{x})$$

$$22.15.3 \quad \lim_{n \rightarrow \infty} \left[\frac{(-1)^n \sqrt{n}}{4^n n!} H_{2n} \left(\frac{x}{2\sqrt{n}} \right) \right] = \frac{1}{\sqrt{\pi}} \cos x$$

$$22.15.4 \quad \lim_{n \rightarrow \infty} \left[\frac{(-1)^n}{4^n n!} H_{2n+1} \left(\frac{x}{2\sqrt{n}} \right) \right] = \frac{2}{\sqrt{\pi}} \sin x$$

$$22.15.5 \quad \lim_{\beta \rightarrow \infty} P_n^{(\alpha, \beta)} \left(1 - \frac{2x}{\beta} \right) = L_n^{(\alpha)}(x)$$

$$22.15.6 \quad \lim_{\alpha \rightarrow \infty} \frac{1}{\alpha^{n/2}} C_n^{(\alpha)} \left(\frac{x}{\sqrt{\alpha}} \right) = \frac{1}{n!} H_n(x)$$

For asymptotic expansions, see [22.5] and [22.17].

22.16. Zeros

For tables of the zeros and associated weight factors necessary for the Gaussian-type quadrature formulas see chapter 25. All the zeros of the orthogonal polynomials are real, simple and located in the interior of the interval of orthogonality.

Explicit and Asymptotic Formulas and Inequalities

Notations:

$x_m^{(n)}$ m th zero of $f_n(x)$ ($x_1^{(n)} < x_2^{(n)} < \dots < x_n^{(n)}$)

$j_{\alpha, m}$, m th positive zero of the Bessel function $J_\alpha(x)$

$\theta_m^{(n)} = \arccos x_{n-m+1}^{(n)}$ ($0 < \theta_1^{(n)} < \theta_2^{(n)} < \dots < \theta_n^{(n)} < \pi$)

$0 < j_{\alpha, 1} < j_{\alpha, 2} < \dots$

	$f_n(x)$	Relation
22.16.1	$P_n^{(\alpha, \beta)}(\cos \theta)$	$\lim_{n \rightarrow \infty} n \theta_m^{(n)} = j_{\alpha, m} \quad (\alpha > -1, \beta > -1)$
22.16.2	$C_n^{(\alpha)}(x)$	$x_m^{(n)} = 1 - \frac{j_{\alpha, m}^2}{2n^2} \left[1 - \frac{2\alpha}{n} + O\left(\frac{1}{n^2}\right) \right]$
22.16.3	$C_n^{(\alpha)}(\cos \theta)$	$\frac{(m+\alpha-1)\pi}{n+\alpha} \leq \theta_m^{(n)} \leq \frac{m\pi}{n+\alpha} \quad (0 \leq \alpha \leq 1)$
22.16.4	$T_n(x)$	$x_m^{(n)} = \cos \frac{2m-1}{2n} \pi$
22.16.5	$U_n(x)$	$x_m^{(n)} = \cos \frac{m}{n+1} \pi$
22.16.6	$P_n(\cos \theta)$	$\begin{cases} \frac{2m-1}{2n+1} \pi \leq \theta_m^{(n)} \leq \frac{2m}{2n+1} \pi \\ \theta_m^{(n)} = \frac{4m-1}{4n+2} \pi + \frac{1}{8n^2} \cot \frac{4m-1}{4n+2} \pi + O(n^{-3}) \end{cases}$
22.16.7	$P_n(x)$	$\begin{cases} x_m^{(n)} = 1 - \frac{j_{\alpha, m}^2}{2n^2} \left[1 - \frac{1}{n} + O(n^{-2}) \right] \\ x_m^{(n)} = 1 - \frac{4\xi_m^{(n)}}{2n+1+\xi_m^{(n)}}; \xi_m^{(n)} = \frac{j_{\alpha, m}^2}{4n+2} \left[1 + \frac{j_{\alpha, m-2}^2}{12(2n+1)^2} \right] + O\left(\frac{1}{n^3}\right) \end{cases}$
22.16.8	$L_n^{(\alpha)}(x)$	$\begin{cases} x_m^{(n)} > \frac{j_{\alpha, m}^2}{4k_n} \\ x_m^{(n)} < \frac{k_m}{k_n} (2k_m + \sqrt{4k_m^2 + \frac{1}{4} - \alpha^2}) \\ x_m^{(n)} = \frac{j_{\alpha, m}^2}{4k_n} \left(1 + \frac{2(\alpha^2-1) + j_{\alpha, m}^2}{48k_n^2} \right) + O(n^{-3}) \end{cases} \quad \left\{ \begin{array}{l} k_r = r + \frac{\alpha+1}{2} \end{array} \right.$

For error estimates see [22.6].

22.17. Orthogonal Polynomials of a Discrete Variable

In this section some polynomials $f_n(x)$ are listed which are orthogonal with respect to the scalar product

$$(f_n, f_m) = \sum_i w^*(x_i) f_n(x_i) f_m(x_i).$$

The x_i are the integers in the interval $a \leq x_i \leq b$ and $w^*(x_i)$ is a positive function such that

$\sum_i w^*(x_i)$ is finite. The constant factor which is still free in each polynomial when only the orthogonality condition is given is defined here by the explicit representation (which corresponds to the Rodrigues' formula)

$$22.17.2 \quad f_n(x) = \frac{1}{r_n w^*(x)} \Delta^n [w^*(x) g(x, n)]$$

where $g(x, n) = g(x)g(x-1) \dots g(x-n+1)$ and $g(x)$ is a polynomial in x independent of n .

Name	a	b	$w^*(x)$	r_n	$g(x, n)$	Remarks
Chebyshev	0	$N-1$	1	$1/n!$	$\binom{x}{n} \binom{x-N}{n}$	
Krawtchouk	0	N	$p^x q^{N-x} \binom{N}{x}$	$(-1)^n n!$	$\frac{q^n x!}{(x-n)!}$	$p, q > 0;$ $p+q=1$
Charlier	0	∞	$\frac{e^{-a} a^x}{x!}$	$(-1)^n \sqrt{a^n n!}$	$\frac{x!}{(x-n)!}$	$a > 0$
Meixner	0	∞	$\frac{c^x \Gamma(b+x)}{\Gamma(b)x!}$	c^n	$\frac{x!}{(x-n)!}$	$b > 0, 0 < c < 1$
Hahn	0	∞	$\frac{\Gamma(b)\Gamma(c+x)\Gamma(d+x)}{x!\Gamma(b+x)\Gamma(c)\Gamma(d)}$	$n!$	$\frac{x!\Gamma(b+x)}{(x-n)!\Gamma(b+x-n)}$	

For a more complete list of the properties of these polynomials see [22.5] and [22.17].

Numerical Methods

22.18. Use and Extension of the Tables

Evaluation of an orthogonal polynomial for which the coefficients are given numerically.

Example 1. Evaluate $L_6(1.5)$ and its first and second derivative using Table 22.10 and the Horner scheme.

1	-36	450	-2400	5400	-4320	720
$x=1.5$	1.5	-51.75	597.375	-2703.9375	4044.09375	-413.859375
1	-34.5	398.25	-1802.625	2696.0625	-275.90625	306.140625
1.5	1.5	-49.5	523.125	-1919.25	1165.21875	$L_6 = \frac{306.140625}{720}$ $= .42519 \ 53$
1	-33.0	348.75	-1279.500	776.8125	889.3125	
1.5	1.5	-47.25	452.250	-1240.875		$L'_6 = \frac{889.3125}{720}$ $= 1.23515 \ 625$
1	-31.5	301.50	-827.250	-464.0625		$L''_6 = 2 \frac{[-464.0625]}{720}$ $= -1.28906 \ 25$

Evaluation of an orthogonal polynomial using the explicit representation when the coefficients are not given numerically.

If an isolated value of the orthogonal polynomial $f_n(x)$ is to be computed, use the proper explicit expression rewritten in the form

$$f_n(x) = d_n(x) a_0(x)$$

and generate $a_0(x)$ recursively, where

$$a_{m-1}(x) = 1 - \frac{b_m}{c_m} f(x) a_m(x) \quad (m=n, n-1, \dots, 2, 1, a_n(x)=1).$$

The $d_n(x)$, b_m , c_m , $f(x)$ for the polynomials of this chapter are listed in the following table:

$f_n(x)$	$d_n(x)$	b_m	c_m	$f(x)$
$P_n^{(\alpha, \beta)}$	$\binom{n+\alpha}{n}$	$(n-m+1)(\alpha+\beta+n+m)$	$2m(\alpha+m)$	$1-x$
$C_{2n}^{(\alpha)}$	$(-1)^n \frac{(\alpha)_n}{n!}$	$2(n-m+1)(\alpha+n+m-1)$	$m(2m-1)$	x^2
$C_{2n+1}^{(\alpha)}$	$(-1)^n \frac{(\alpha)_{n+1}}{n!} 2x$	$2(n-m+1)(\alpha+n+m)$	$m(2m+1)$	x^2
T_{2n}	$(-1)^n$	$2(n-m+1)(n+m-1)$	$m(2m-1)$	x^2
T_{2n+1}	$(-1)^n (2n+1)x$	$2(n-m+1)(n+m)$	$m(2m+1)$	x^2
U_{2n}	$(-1)^n$	$2(n-m+1)(n+m)$	$m(2m-1)$	x^2
U_{2n+1}	$(-1)^n (2n+1)x$	$2(n-m+1)(n+m+1)$	$m(2m+1)$	x^2
P_{2n}	$\frac{(-1)^n}{4^n} \binom{2n}{n}$	$(n-m+1)(2n+2m-1)$	$m(2m-1)$	x^2
P_{2n+1}	$\frac{(-1)^n}{4^n} \binom{2n+1}{n} (n+1)x$	$(n-m+1)(2n+2m+1)$	$m(2m+1)$	x^2
$L_n^{(\alpha)}$	$\binom{n+\alpha}{n}$	$n-m+1$	$m(\alpha+m)$	x
H_{2n}	$(-1)^n \frac{(2n)!}{n!}$	$2(n-m+1)$	$m(2m-1)$	x^2
H_{2n+1}	$(-1)^n \frac{(2n+1)!}{n!} 2x$	$2(n-m+1)$	$m(2m+1)$	x^2

Example 2. Compute $P_8^{(1/2, 3/2)}(2)$. Here $d_8 = \binom{8.5}{8} = 3.33847$, $f(2) = -1$.

m	8	7	6	5	4	3	2	1	0
a_m	1	1.132353	1.366667	1.841026	3.008392	6.849651	26.44156	223.1091	6545.533
b_m	18	34	48	60	70	78	84	88	90
c_m	136	105	78	55	36	21	10	3	0

$$P_8^{(1/2, 3/2)}(2) = d_8 a_0(2) = (3.33847)(6545.533) = 21852.07$$

Evaluation of orthogonal polynomials by means of their recurrence relations

Example 3. Compute $C_n^{(1)}(2.5)$ for $n=2, 3, 4, 5, 6$.

From Table 22.2 $C_0^{(1)}=1$, $C_1^{(1)}=1.25$ and from 22.7 the recurrence relation is

$$C_{n+1}^{(1)}(2.5) = [5(n+\frac{1}{2})C_n^{(1)}(2.5) - (n-\frac{1}{2})C_{n-1}^{(1)}(2.5)] \frac{1}{n+1}.$$

n	2	3	4	5	6
$C_n^{(1)}(2.5)$	3.65625	13.08594	50.87648	207.0649	867.7516

Check: Compute $C_6^{(1)}(2.5)$ by the method of Example 2.

Change of Interval of Orthogonality

In some applications it is more convenient to use polynomials orthogonal on the interval $[0, 1]$. One can obtain the new polynomials from the ones given in this chapter by the substitution $x=2\bar{x}-1$. The coefficients of the new polynomial can be computed from the old by the following recursive scheme, provided the standardization is not changed. If

$$f_n(x) = \sum_{m=0}^n a_m x^m, \quad f_n^*(x) = f_n(2x-1) = \sum_{m=0}^n a_m^* x^m$$

then the a_m^* are given recursively by the a_m through the relations

$$a_m^{(j)} = 2a_m^{(j-1)} - a_{m+1}^{(j-1)}; \quad m=n-1, n-2, \dots, j; \quad j=0, 1, 2, \dots, n$$

$$a_m^{(-1)} = a_m/2, \quad m=0, 1, 2, \dots, n$$

$$a_n^{(j)} = 2^j a_n, \quad j=0, 1, 2, \dots, n \text{ and } a_m^{(n)} = a_m^*; \quad m=0, 1, 2, \dots, n.$$

Example 4. Given $T_5(x) = 5x - 20x^3 + 16x^5$, find $T_5^*(x)$.

$m \backslash j$	5	4	3	2	1	0
-1	$8 = a_5^{(-1)}$	0	$-10 = a_4^{(-1)}$	0	$2.5 = a_3^{(-1)}$	0
0	16	-16	-4	4	1	$-1 = a_0^*$
1	32	-64	56	-48	$50 = a_1^*$	
2	64	-192	304	-400	a_2^*	
3	128	-512	1120	a_3^*		
4	256	-1280	a_4^*			
5	$512 = a_5^*$					

Hence, $T_5^*(x) = 512x^5 - 1280x^4 + 1120x^3 - 400x^2 + 50x - 1$.

22.19. Least Square Approximations

Problem: Given a function $f(x)$ (analytically or in form of a table) in a domain D (which may be a continuous interval or a set of discrete points).^{*} Approximate $f(x)$ by a polynomial $F_n(x)$ of given degree n such that a weighted sum of the squares of the errors in D is least.

Solution: Let $w(x) \geq 0$ be the weight function chosen according to the relative importance of the errors in different parts of D . Let $f_m(x)$ be orthogonal polynomials in D relative to $w(x)$, i.e. $(f_m, f_n) = 0$ for $m \neq n$, where

$$(f, g) = \begin{cases} \int_D w(x) f(x) g(x) dx & \text{if } D \text{ is a continuous interval} \\ \sum_{m=1}^N w(x_m) f(x_m) g(x_m) & \text{if } D \text{ is a set of } N \text{ discrete points } x_m. \end{cases}$$

Then

$$F_n(x) = \sum_{m=0}^n a_m f_m(x)$$

where

$$a_m = (f, f_m) / (f_m, f_m).$$

^{*} $f(x)$ has to be square integrable, see e.g. [22.17].

D a Continuous Interval

Example 5. Find a least square polynomial of degree 5 for $f(x) = \frac{1}{1+x}$, in the interval $2 \leq x \leq 5$, using the weight function

$$w(x) = \frac{1}{\sqrt{(x-2)(5-x)}}$$

which stresses the importance of the errors at the ends of the interval.

Reduction to interval $[-1, 1]$, $t = \frac{2x-7}{3}$

$$w(x(t)) = \frac{2}{3} \frac{1}{\sqrt{1-t^2}}$$

From 22.2, $f_m(t) = T_m(t)$ and

$$a_m = \frac{4}{3\pi} \int_{-1}^1 \frac{1}{\sqrt{1-t^2}} \frac{1}{t+3} T_m(t) dt \quad (m \neq 0)$$

$$a_0 = \frac{2}{3\pi} \int_{-1}^1 \frac{1}{\sqrt{1-t^2}} \frac{dt}{t+3}$$

Evaluating the integrals numerically we get

$$\frac{1}{1+x} \sim .235703 - .080880T_1\left(\frac{2x-7}{3}\right) + .013876T_2\left(\frac{2x-7}{3}\right) - .002380T_3\left(\frac{2x-7}{3}\right) \\ + .000408T_4\left(\frac{2x-7}{3}\right) - .000070T_5\left(\frac{2x-7}{3}\right)$$

D a Set of Discrete Points

If $x_m = m (m=0, 1, 2, \dots, N)$ and $w(x)=1$, use the Chebyshev polynomials in the discrete range 22.17. It is convenient to introduce here a slightly different standardization such that

$$f_n(x) = \sum_{m=0}^n (-1)^m \binom{n}{m} \binom{n+m}{m} \frac{x!(N-m)!}{(x-m)!N!} \\ (f_n, f_n) = \frac{(N+n+1)!(N-n)!}{(2n+1)(N!)^2}$$

Recurrence relation: $f_0(x)=1, f_1(x)=1-\frac{2x}{N}$

$$(n+1)(N-n)f_{n+1}(x) = (2n+1)(N-2x)f_n(x) - n(N+n+1)f_{n-1}(x)$$

Example 6. Approximate in the least square sense the function $f(x)$ given in the following table by a third degree polynomial.

x	$f(x)$	$\bar{x} = \frac{x-10}{2}$	$f_0(\bar{x})$	$f_1(\bar{x})$	$f_2(\bar{x})$	$f_3(\bar{x})$
10	.3162	0	1	1	1	1
12	.2887	1	1	1/2	-1/2	-2
14	.2673	2	1	0	-1	0
16	.2500	3	1	-1/2	-1/2	2
18	.2357	4	1	-1	1	-1

$$f_0(\bar{x}) \quad f_1(\bar{x}) \quad f_2(\bar{x}) \quad f_3(\bar{x})$$

$$(f_n, f_n) = \sum_{\bar{x}=0}^4 f_n^2(\bar{x})$$

$$5 \quad 2.5 \quad 3.5 \quad 10$$

$$(f, f_n) = \sum_{\bar{x}=0}^4 f_n(\bar{x})f(2\bar{x}+10)$$

$$1.3579 \quad .09985 \quad .01525 \quad .0031$$

$$a_n = \frac{(f, f_n)}{(f_n, f_n)}$$

$$.271580 \quad .039940 \quad .0043571 \quad .000310$$

$$f(x) \sim .27158 + .03994(3.5 - .25x) + .0043571(23.5 - 3.5x + .125x^2) + .00031(266 - 59.8333x \\ + 4.375x^2 - .10417x^3)$$

$$f(x) \sim .59447 - .043658x + .0019009x^2 - .000032292x^3$$

22.20. Economization of Series

Problem: Given $f(x) = \sum_{m=0}^n a_m x^m$ in the interval $-1 \leq x \leq 1$ and $R > 0$. Find $\bar{f}(x) = \sum_{m=0}^N b_m x^m$ with N as small as possible, such that $|\bar{f}(x) - f(x)| < R$.

Solution: Express $f(x)$ in terms of Chebyshev polynomials using Table 22.3,

$$f(x) = \sum_{m=0}^n b_m T_m(x)$$

Then, since $|T_m(x)| \leq 1 (-1 \leq x \leq 1)$

$$\bar{f}(x) = \sum_{m=0}^N b_m T_m(x)$$

within the desired accuracy if

$$\sum_{m=N+1}^n |b_m| < R$$

$\bar{f}(x)$ is evaluated most conveniently by using the recurrence relation (see 22.7).

Example 7. Economize $f(x)=1+x/2+x^2/3+x^3/4+x^4/5+x^5/6$ with $R=.05$.

From Table 22.3

$$f(x) = \frac{1}{120} [149T_0(x) + 32T_2(x) + 3T_4(x)] + \frac{1}{96} [76T_1(x) + 11T_3(x) + T_5(x)]$$

so

$$\bar{f}(x) = \frac{1}{120} [149T_0(x) + 32T_2(x)] + \frac{1}{96} [76T_1(x) + 11T_3(x)]$$

since

$$|\bar{f}(x) - f(x)| \leq \frac{1}{40} + \frac{1}{96} < .05$$

References

Texts

- [22.1] Bibliography on orthogonal polynomials, Bull. of the National Research Council No. 103, Washington, D.C. (1940).
- [22.2] P. L. Chebyshev, Sur l'interpolation. Oeuvres, vol. 2, pp. 59-68.
- [22.3] R. Courant and D. Hilbert, Methods of mathematical physics, vol. 1, ch. 7 (Interscience Publishers, New York, N.Y., 1953).
- [22.4] G. Doetsch, Die in der Statistik seltener Ereignisse auftretenden Charlierschen Polynome und eine damit zusammenhängende Differential-differenzengleichung, Math. Ann. **109**, 257-266 (1934).
- [22.5] A. Erdélyi et al., Higher transcendental functions, vol. 2, ch. 10 (McGraw-Hill Book Co., Inc., New York, N.Y., 1953).
- [22.6] L. Gatteschi, Limitazione degli errori nelle formule asintotiche per le funzioni speciali, Rend. Sem. Mat. Univ. Torino **16**, 83-94 (1956-57).
- [22.7] T. L. Geronimus, Teoria ortogonalnikh mnogochlenov (Moscow, U.S.S.R., 1950).
- [22.8] W. Hahn, Über Orthogonalpolynome, die q-Differenzengleichungen genügen, Math. Nachr. **2**, 4-34 (1949).
- [22.9] St. Kaczmarz and H. Steinhaus, Theorie der Orthogonalreihen, ch. 4 (Chelsea Publishing Co., New York, N.Y., 1951).
- [22.10] M. Krawtchouk, Sur une généralisation des polynomes d'Hermite, C.R. Acad. Sci. Paris **187**, 620-622 (1929).
- [22.11] C. Lanczos, Trigonometric interpolation of empirical and analytical functions, J. Math. Phys. **17**, 123-199 (1938).
- [22.12] C. Lanczos, Applied analysis (Prentice-Hall, Inc., Englewood Cliffs, N.J., 1956).
- [22.13] W. Magnus and F. Oberhettinger, Formeln und Sätze für die speziellen Funktionen der mathematischen Physik, ch. 5, 2d ed. (Springer-Verlag, Berlin, Germany, 1948).

- [22.14] J. Meixner, Orthogonale Polynomsysteme mit einer besonderen Gestalt der erzeugenden Funktion, J. London Math. Soc. **9**, 6-13 (1934).
- [22.15] G. Sansone, Orthogonal functions, Pure and Applied Mathematics, vol. IX (Interscience Publishers, New York, N.Y., 1959).
- [22.16] J. Shohat, Théorie générale des polynômes orthogonaux de Tchebichef, Mém. Soc. Math. **66** (Gauthier-Villars, Paris, France, 1934).
- [22.17] G. Szegő, Orthogonal polynomials, Amer. Math. Soc. Colloquium Publications **23**, rev. ed. (1959).
- [22.18] F. G. Tricomi, Vorlesungen über Orthogonalreihen, chs. 4, 5, 6 (Springer-Verlag, Berlin, Germany, 1955).

Tables

- [22.19] British Association for the Advancement of Science, Legendre Polynomials, Mathematical Tables, Part vol. A (Cambridge Univ. Press, Cambridge, England, 1946). $P_n(x)$, $x=0(.01)6$, $n=1(1)12$, 7-8D.
- [22.20] N. R. Jörgensen, Undersøgelser over frekvensflader og korrelation (Busck, Copenhagen, Denmark, 1916). $He_n(x)$, $x=0(.01)4$, $n=1(1)6$, exact.
- [22.21] L. N. Karmazina, Tablitsy polinomov Jacobi (Izdat. Akad. Nauk SSSR., Moscow, U.S.S.R., 1954). $G_n(p, q, x)$, $x=0(.01)1$, $q=.1(1)1$, $p=1.1(1)3$, $n=1(1)5$, 7D.
- [22.22] National Bureau of Standards, Tables of Chebyshev polynomials $S_n(x)$ and $C_n(x)$, Applied Math. Series 9 (U.S. Government Printing Office, Washington, D.C., 1952). $x=0(.001)2$, $n=2(1)12$, 12D; Coefficients for $T_n(x)$, $U_n(x)$, $C_n(x)$, $S_n(x)$ for $n=0(1)12$.
- [22.23] J. B. Russel, A table of Hermite functions, J. Math. Phys. **12**, 291-297 (1933). $e^{-x^2/2}H_n(x)$, $x=0(.04)1(.1)4(.2)7(.5)8$, $n=0(1)11$, 5D.
- [22.24] N. Wiener, Extrapolation, interpolation and smoothing of stationary time series (John Wiley & Sons, Inc., New York, N.Y., 1949). $L_n(x)$, $n=0(1)5$, $x=0(.01).1(.1)18(.2)20(.5)21(1)26(2)30$, 3-5D.